

Strong meager properties for filters

Claude Laflamme*

Department of Mathematics and Statistics
University of Calgary
Calgary, Alberta
Canada T2N 1N4

Abstract

We analyze several “strong meager” properties for filters on the natural numbers between the classical Baire property and a filter being F_σ . Two such properties have been studied by Talagrand and a few more combinatorial ones are investigated. In particular, we define the notion of a P^+ -filter, a generalization of the traditional concept of P -filter, and prove the existence of a non-meager P^+ -filter. Our motivation lies in understanding the structure of filters generated by complements of members of a maximal almost disjoint family.

1 Introduction

We investigate several “smallness” properties of filters as strengthening of the meager property. Indeed, although a meager filter might be thought of as small, it may happen for example that adding one set to a meager filter yields an ultrafilter (see the section on examples). Our motivation lies in the unresolved relationship between the minimum size of a maximal almost disjoint (mad) family and the minimum size of a dominating family; the relation with the minimum size of an unbounded family has been settled by Shelah in [4]. Part of the problem lies in the technical question whether filters generated by the complement of members of a mad family can be diagonalized in a forcing extension without adding an unbounded real. This loosely means that we must understand the different possibilities for a filter to code an unbounded function when diagonalized. Very partial results were obtained in [3] where it became apparent that strong meager properties are required on filters for this program to succeed. It is the goal of this paper to analyze some of these stronger meager properties.

*This research was partially supported by NSERC of Canada.
1980 *AMS Subj. Class. (1985 Revision)* Primary 04A20; Secondary 03E05, 03E15, 03E35.
Key words and phrases. Filter, meager, Baire property.

Our terminology is standard but we review the main concepts and notation. The natural numbers will be denoted by ω , ${}^\omega 2$ and ${}^\omega \omega$ denote the collection of functions from ω to 2 and to ω respectively; similarly, $\mathcal{O}(\omega)$ and $[\omega]^\omega$ denote the collection of all and infinite subsets respectively. Two orderings are standard, the first one for subsets of ω is $A \subseteq^* B$ if $A \setminus B$ is finite, the second one for functions is $f \leq^* g$ if $f(n) \leq g(n)$ for all but finitely many n . We can view members of $\mathcal{O}(\omega)$ as members of ${}^\omega 2$ by considering their characteristic functions. We equip ${}^\omega 2$ with the product topology of the discrete space $\{0, 1\}$. A basic neighbourhood is then given by sets of the form

$$\mathcal{O}_s = \{f \in {}^\omega 2 : s \subseteq f\}$$

where $s \in {}^{<\omega} 2$, the collection of finite binary sequences. The terms “nowhere dense”, “meager”, “Baire property” all refer to this topology.

A filter is a collection of subsets of ω closed under finite intersections, supersets and containing all cofinite sets; it is called proper if it contains only infinite sets. Given a collection $\mathcal{X} \subseteq \mathcal{O}(\omega)$, we let $\langle \mathcal{X} \rangle$ denote the filter generated by $\mathcal{X}$. For a filter $\mathcal{F}$, $\mathcal{F}^+$ denotes the collection of all sets X such that $\langle \mathcal{F}, X \rangle$ is a proper filter; it is useful to notice that $X \in \mathcal{F}^+$ if and only if $X^c \notin \mathcal{F}$. We say that $X \in [\omega]^\omega$ diagonalizes a filter $\mathcal{F}$ if $X \subseteq^* Y$ for all $Y \in \mathcal{F}$. The *Fréchet* filter is the collection of cofinite sets, denoted by Fr . Given two filters $\mathcal{F}$ and $\mathcal{G}$, we form a third filter by

$$\mathcal{F} \otimes \mathcal{G} = \{X \subseteq \omega \times \omega : \{n : \{m : (n, m) \in X\} \in \mathcal{G}\} \in \mathcal{F}\}.$$

This filter can be viewed as a filter on ω if desired by fixing a bijection between $\omega \times \omega$ and ω .

The following important result characterizes meager filters in terms of combinatorial properties.

Proposition 1.1 (*Talagrand ([6])*) *The following are equivalent for a filter $\mathcal{F}$:*

1. $\mathcal{F}$ has the Baire property.
2. $\mathcal{F}$ is meager.
3. There is a sequence $n_0 < n_1 < \dots$ such that

$$(\forall X \in \mathcal{F})(\forall^\infty k) X \cap [n_k, n_{k+1}) \neq \emptyset.$$

A family $\mathcal{F} \subseteq {}^\omega \omega$ is called bounded if

$$(\exists g \in {}^\omega \omega)(\forall f \in \mathcal{F}) f \leq^* g.$$

Similarly, a family $\mathcal{F} \subseteq {}^\omega \omega$ is called dominating if

$$(\forall g \in {}^\omega \omega)(\exists f \in \mathcal{F}) g \leq^* f.$$


The cardinals $\mathfrak{b}$ and $\mathfrak{d}$ denote respectively the minimum cardinality of an unbounded or dominating family.

2 Strong meager properties

Definition 2.1 Let $\mathcal{F}$ be a filter on ω .

1. $\mathcal{F}$ is called *completely meager* (Talagrand, [7]) if the filter $\langle \mathcal{F}, X \rangle$ is meager whenever $X^c \notin \mathcal{F}$.
2. $\mathcal{F}$ is called *hereditarily meager* if for each $f \in {}^\omega\omega$, the filter $f(\mathcal{F})$ is meager (or improper).
3. $\mathcal{F}$ is called *weakly hereditarily meager* if for each $f \in {}^\omega\omega$, either $f^{-1}(i)$ is compatible with $\mathcal{F}$ for some i , or else $f(\mathcal{F})$ is meager.
4. $\mathcal{F}$ has the *strong Baire property* (Talagrand, [7]) if $\mathcal{F} \cap \mathcal{C}$ has the Baire property relative to $\mathcal{C}$ for each closed $\mathcal{C} \subseteq \wp(\omega)$.
5. $\mathcal{F}$ is a P^+ -filter if given any decreasing sequence $\langle X_0 \supseteq X_1 \supseteq X_2 \supseteq \dots \rangle$ from $\mathcal{F}^+$, there is an $X \in \mathcal{F}^+$ such that $X \subseteq^* X_n$ for each n .
6. $\mathcal{F}$ is a *strong P^+ -filter* if given any sequence $\langle X_0, X_1, X_2, \dots \rangle$ from $\mathcal{F}^+$, there is an $f \in {}^\omega\omega$ such that for all $X \in \mathcal{F}$, $X \cap X_n \cap f(n) \neq \emptyset$ for all but finitely many n .

We shall prove the following implications between these notions. In a subsequent section, we shall show by examples that many of these implications do not reverse. Some open questions remain.



The following lemmas will only reformulate these definitions into more uniform and manageable ones; the proofs of the first three are left to the reader.

Lemma 2.2 *The following are equivalent for a given filter $\mathcal{F}$:*

1. $\mathcal{F}$ is weakly hereditarily meager.
2. For each sequence $\langle X_i : i \in \omega \rangle$ such that $\bigcup_i X_i \in \mathcal{F}$ and each $X_i^c \in \mathcal{F}$,

$$(\exists n_0 < n_1 < \dots)(\forall Y \in \mathcal{F})(\forall^\infty k) Y \cap \bigcup_{i \in [n_k, n_{k+1})} X_i \neq \emptyset.$$

Lemma 2.3 *The following are equivalent for a given filter $\mathcal{F}$:*

1. $\mathcal{F}$ is completely meager.
2. For each sequence of finite sets $\langle X_i : i \in \omega \rangle$ such that $\bigcup_i X_i \in \mathcal{F}^+$,

$$(\exists n_0 < n_1 < \dots)(\forall Y \in \mathcal{F})(\forall^\infty k) Y \cap \bigcup_{i \in [n_k, n_{k+1})} X_i \neq \emptyset.$$

Lemma 2.4 *The following are equivalent for a given filter $\mathcal{F}$:*

1. $\mathcal{F}$ is hereditarily meager.
2. For each sequence $\langle X_i : i \in \omega \rangle$ such that $\bigcup_i X_i \in \mathcal{F}^+$ and each $X_i^c \in \mathcal{F}$,

$$(\exists n_0 < n_1 < \dots)(\forall Y \in \mathcal{F})(\forall^\infty k) Y \cap \bigcup_{i \in [n_k, n_{k+1})} X_i \neq \emptyset.$$

3. For each sequence $\langle X_i : i \in \omega \rangle$ such that $\bigcup_i X_i \in \mathcal{F}$, $\bigcup_{i < k} X_i \notin \mathcal{F}$ for each k and only finitely many X_i belong to $\mathcal{F}^+$,

$$(\exists n_0 < n_1 < \dots)(\forall Y \in \mathcal{F})(\forall^\infty k) Y \cap \bigcup_{i \in [n_k, n_{k+1})} X_i \neq \emptyset.$$

Lemma 2.5 *The following are equivalent for a given filter $\mathcal{F}$:*

1. $\mathcal{F}$ is a completely meager P^+ -filter.
2. For each sequence $\langle X_i : i \in \omega \rangle$ such that $\bigcup_{i \geq n} X_i \in \mathcal{F}^+$ for each n ,

$$(\exists h \in {}^\omega \omega)(\exists n_0 < n_1 < \dots)(\forall Y \in \mathcal{F})(\forall^\infty k) Y \cap \bigcup_{i \in [n_k, n_{k+1})} X_i \cap h(i) \neq \emptyset.$$

3. For each sequence $\langle X_i : i \in \omega \rangle \subseteq \mathcal{F}^+$,

$$(\exists h \in {}^\omega \omega)(\exists n_0 < n_1 < \dots)(\forall Y \in \mathcal{F})(\forall^\infty k) Y \cap \bigcup_{i \in [n_k, n_{k+1})} X_i \cap h(i) \neq \emptyset.$$

Proof: 1 $\rightarrow$ 2: Let $\mathcal{F}$ be a completely meager P^+ -filter and fix a sequence $\langle X_i : i \in \omega \rangle$ such that $\bigcup_{i \geq n} X_i \in \mathcal{F}^+$ for each n . Define $Y_n = \bigcup_{i \geq n} X_i$. Then $Y_0 \supseteq Y_1 \supseteq \dots$ is a decreasing sequence from $\mathcal{F}^+$ and therefore there is an $X \in \mathcal{F}^+$ almost included in each Y_i . Since $\langle \mathcal{F}, X \rangle$ must be meager, choose $m_0 < m_1 < \dots$ such that

$$(\forall Y \in \mathcal{F})(\forall^\infty k) Y \cap X \cap [m_k, m_{k+1}) \neq \emptyset.$$

Now choose $n_0 < n_1 < \dots$ and $h \in {}^\omega \omega$ such that

$$(\forall k)(\exists \ell) [m_\ell, m_{\ell+1}) \cap X \subseteq \bigcup_{i \in [n_k, n_{k+1})} X_i \cap h(i).$$

2 $\rightarrow$ 3 is immediate.

3 $\rightarrow$ 1: To show that $\mathcal{F}$ is a P^+ -filter, fix a decreasing sequence $X_0 \supseteq X_1 \supseteq \dots$ from $\mathcal{F}^+$. By assumption, there are $h \in {}^\omega \omega$ and a sequence $n_0 < n_1 < \dots$ such that

$$(\forall Y \in \mathcal{F}) Y \cap \bigcup_{i \in [n_k, n_{k+1})} X_i \cap h(i) \neq \emptyset.$$

If we let $X = \bigcup_k X_{n_k} \cap h(n_k) = \bigcup_i X_i \cap h(i)$, then $X \in \mathcal{F}^+$ and is almost included in each X_i .

To show that $\mathcal{F}$ is completely meager, let $X \in \mathcal{F}^+$ be given and use the assumptions on $X_i = X \setminus i$. $\square$

Now we are ready to prove the diagram implications; in view of the above lemmas, all are straightforward except possibly two of them which we prove in detail.

Proposition 2.6 *A strong P^+ -filter has the strong Baire property.*

Proof: We actually show that a strong P^+ -filter has the Baire property relative to any set $\mathcal{X} \subseteq {}^\omega 2$. Indeed, let

$$\mathcal{T} = \{s \in {}^{<\omega} 2 : \mathcal{O}_s \cap \mathcal{X} \subseteq \mathcal{F}\},$$

and put $\mathcal{O} = \bigcup_{s \in \mathcal{T}} \mathcal{O}_s$. We show that $(\mathcal{F} \cap \mathcal{X}) \Delta (\mathcal{O} \cap \mathcal{X})$ is meager in $\mathcal{X}$. For this, let $\mathcal{S} = \{s \in {}^{<\omega} 2 : \mathcal{O}_s \cap \mathcal{X} \neq \emptyset \text{ and } \mathcal{O}_s \cap \mathcal{X} \not\subseteq \mathcal{F}\}$; enumerate $\mathcal{S}$ as $\langle s_n : n \in \omega \rangle$. For each n , choose $A_n \in (\mathcal{O}_{s_n} \cap \mathcal{X}) \setminus \mathcal{F}$ and put $X_n = A_n^c \setminus lh(s) \in \mathcal{F}^+$.

By assumption, there is an $h \in {}^\omega\omega$ such that for all $Y \in \mathcal{F}$, $Y \cap X_n \cap h(n) \neq \emptyset$ for all but finitely many n . Define now

$$M_k = \{X : (\forall \ell \geq k) X \cap X_\ell \neq \emptyset\} \setminus \mathcal{O}.$$

Clearly $(\mathcal{F} \setminus \mathcal{O}) \subseteq \bigcup_k M_k$ and each M_k is nowhere dense in $\mathcal{X}$. $\square$

Proposition 2.7 *Every F_σ -filter is a strong P^+ -filter.*

Proof: Let $\mathcal{F} = \bigcup_n \mathcal{C}_n$ be a countable increasing union of closed sets and $\langle X_n : n \in \omega \rangle \subseteq \mathcal{F}^+$. Put $\mathcal{C} = \bigcup \{X \cup n : X \in \mathcal{C}_n\}$ and define $\mathcal{C}'_n = \mathcal{C} \cup X_n$. Choose $h \in {}^\omega\omega$ such that for all n ,

$$(\forall Y, Z \in \mathcal{C}'_n) Y \cap Z \cap [n, h(n)) \neq \emptyset.$$

The existence of such a function is guaranteed by the fact that $\mathcal{C}'_n$ is closed. Now if $Y \in \mathcal{F}$, say $Y \in \mathcal{C}_i$, then $Y \cup i \in \mathcal{C}$ and therefore $(Y \cup i) \cap X_n \cap h(n) \neq \emptyset$ for all n , and thus $Y \cap X_n \cap h(n) \neq \emptyset$ for all $n \geq i$ as desired. $\square$

3 Examples

We now produce examples showing that many implications in the diagram are not reversible.

Example 3.1 *A weakly hereditarily meager filter which is not completely meager.*

Let $\mathcal{U}$ be any ultrafilter not containing say the even numbers E . Define $\mathcal{F} = \{E \cup X : X \in \mathcal{U}\}$. Then $\langle \mathcal{F}, E^c \rangle = \mathcal{U}$ is not meager and thus $\mathcal{F}$ is not completely meager.

Now let $\langle X_i : i \in \omega \rangle$ be given such that $\bigcup_i X_i \in \mathcal{F}$ and each $X_i^c \in \mathcal{F}$. Choose a sequence $n_0 < n_1 < \dots$ such that for each k , $(E \setminus k) \cap \bigcup_{i \in [n_k, n_{k+1})} X_i \neq \emptyset$. This is possible as $E \cap \bigcup_{i < k} X_i$ is finite for each k . Since any $Y \in \mathcal{F}$ almost contains E , we conclude that $\mathcal{F}$ is weakly hereditarily meager by lemma 2.2. $\square$

Example 3.2 *A completely meager filter which is not weakly hereditarily meager.*

This example comes essentially from [7], where a completely meager filter without the strong Baire property is produced. Let again $\mathcal{U}$ be an ultrafilter and let $\mathcal{F} = \mathcal{U} \otimes Fr$.

Claim 1: $\mathcal{F}$ is completely meager.

Indeed let $X \subseteq \omega \times \omega$ be compatible with $\mathcal{F}$; this means that $X_n = \{(n, m) : m \in X\}$ is infinite for infinitely many n (actually on a set in $\mathcal{U}$). Define a sequence of finite sets $\langle s_i : i \in \omega \rangle$ such that $(s_{i+1} \setminus s_i) \cap X_j \neq \emptyset$ for each $j \leq i$ such that X_j is infinite. Clearly

$$(\forall Y \in \mathcal{F})(\forall^\infty i) Y \cap X \cap s_i \neq \emptyset$$

and thus $\langle \mathcal{F}, X \rangle$ is meager.

Claim 2: $\mathcal{F}$ is not weakly hereditarily meager.

Define $f \in {}^\omega\omega$ by $f(n, m) = n$, the downward projection. Then no $f^{-1}(i)$ is compatible with $\mathcal{F}$ but $f(\mathcal{F}) = \mathcal{U}$ is not meager. $\square$

Example 3.3 *A completely meager P^+ -filter which is not a strong P^+ -filter.*

We build a mad family $\mathcal{A}$ such that the filter $\mathcal{F}(\mathcal{A})$ has the required properties. It is essentially proved in [3] that any filter of the form $\mathcal{F}(\mathcal{A})$ will necessarily be a completely meager P^+ -filter; we include the proof for completeness.

CLAIM 1: All filters of the form $\mathcal{F}(\mathcal{A})$ are completely meager P^+ -filters.

Notice that a set $X \in \mathcal{F}(\mathcal{A})^+$ if and only if X has infinite intersection with infinitely many members of $\mathcal{A}$. Now if $Y_0 \supseteq Y_1 \supseteq \dots$ are from $\mathcal{F}(\mathcal{A})^+$, choose $\{A_n : n \in \omega\} \subseteq \mathcal{A}$ such $Y_n \cap A_m \neq \emptyset$ for each n, m ; finally define a sequence $\langle n_k : k \in \omega \rangle$ such that

$$Y_k \cap A_\ell \cap [n_k, n_{k+1}) \neq \emptyset$$

for each k and $\ell \leq k$. Clearly $Y = \bigcup_k Y_k \cap [n_k, n_{k+1}) \in \mathcal{F}(\mathcal{A})^+$ and is almost included in each Y_n . Actually we see that for any $X \in \mathcal{F}(\mathcal{A})$, $X \cap Y \cap [n_k, n_{k+1}) \neq \emptyset$ for all but finitely many k and thus we have also proved that $\mathcal{F}(\mathcal{A})$ is completely meager.

Now to make sure that $\mathcal{F}(\mathcal{A})$ is not a strong P^+ -filter, first fix an almost disjoint family $\mathcal{B} = \{B_\alpha : \alpha < \mathfrak{c}\}$ of size continuum and fix an enumeration of ${}^\omega\omega = \{f_\alpha : \alpha < \mathfrak{c}\}$. Fix further a partition $\langle X_n : n \in \omega \rangle$ of ω . Now we define an almost disjoint family

$$\mathcal{A}' = \left\{ \bigcup_{n \in B_\alpha} X_n \cap f_\alpha(n) : \alpha < \mathfrak{c} \right\}.$$

Let $\mathcal{A}$ be any mad family extending $\mathcal{A}'$ together with a partition of the X_n 's into countably many infinite sets (to make sure that $X_n \in \mathcal{F}(\mathcal{A})^+$). Clearly $\mathcal{F}(\mathcal{A})$ is not a strong P^+ -filter.

Example 3.4 *A filter with the strong Baire property which is not a P^+ -filter.*

Let $\mathcal{F} = Fr \otimes Fr$. Then $\mathcal{F}$ is a Borel filter and thus has the strong Baire property. If we define $Y_n = \{(\ell, m) : \ell \geq n\} \in \mathcal{F}$, then $\mathcal{F}$ contains the complement of all Y almost included in each Y_n .

Example 3.5 *A non-meager P^+ -filter.*

It is worth noticing that the concept of P^+ -filter and the traditional notion of P-filter are incomparable, that is one does not imply the other. The difficult question of producing a non-meager P-filter has been around for a some time, see [2] for some results and applications in this area.

We show that Simon's example ([5]) of a non-meager filter generated by $\mathfrak{b}$ sets is actually a P^+ -filter. As in his construction, we consider two possibilities.

Case 1: ($\mathfrak{b} < \mathfrak{d}$) Choose an unbounded family $\langle f_\alpha : \alpha < \mathfrak{b} \rangle$ of strictly increasing functions such that

$$\alpha < \beta \rightarrow f_\alpha <^* f_\beta.$$

Fix now $g \in {}^\omega\omega$ not dominated by any f_α , put

$$X_\alpha = \{n : f_\alpha(n) < g(n)\}$$

and define $\mathcal{F} = \langle X_\alpha : \alpha < \mathfrak{b} \rangle$. $\mathcal{F}$ is a P-filter and the next lemma shows that it really is a P^+ -filter.

Lemma 3.6 *Any filter generated by less than $\mathfrak{d}$ sets is a P^+ -filter.*

Let $\mathcal{H}$ be a filter generated by $\langle X_\alpha : \alpha < \lambda \rangle$ for some $\lambda < \mathfrak{d}$, and fix a descending sequence $Y_0 \supseteq Y_1 \supseteq \dots$ from $\mathcal{H}^+$. For each $\alpha < \lambda$, choose $h_\alpha \in {}^\omega\omega$ such that

$$(\forall n)(Y_n \cap X_\alpha \cap [n, h_\alpha(n)) \neq \emptyset.$$

Since $\lambda < \mathfrak{d}$, fix $h \in {}^\omega\omega$ not dominated by any h_α and let $Y = \bigcup_n Y_n \cap h(n)$ almost included in each Y_n . But for each α , if $h_\alpha(n) \leq h(n)$, $Y \cap X_\alpha \cap [n, h(n)) \neq \emptyset$ and therefore $Y \in \mathcal{H}^+$. $\square$

Finally we must show that $\mathcal{F}$ is non-meager. So suppose we are given a sequence $n_0 < n_1 < \dots$ such that

$$(\forall X \in \mathcal{F})(\forall^\infty k) X \cap [n_k, n_{k+1}) \neq \emptyset.$$

Then for each α , choose m such that $X_\alpha \cap [n_k, n_{k+1}) \neq \emptyset$ for each $k \geq m$. Therefore for each ℓ , $f_\alpha(\ell) \leq n_{\ell+m+1}$ and thus for all $\ell \geq m+1$ we have $f_\alpha(\ell) \leq n_{2\ell}$. We conclude that the family $\langle f_\alpha : \alpha < \mathfrak{b} \rangle$ is bounded by $f(\ell) = n_{2\ell}$, a contradiction.

Case 2: ($\mathfrak{b} = \mathfrak{d}$) Choose a scale $\langle f_\alpha : \alpha < \mathfrak{b} = \mathfrak{d} \rangle$ such that

1. Each f_α is strictly increasing and positive.
2. $(\forall \beta < \alpha)(\forall^\infty n)(\exists \ell) f_\alpha(n) < f_\beta(\ell) < f_\beta(\ell+1) < f_\alpha(n+1)$.

For each α , put $X_\alpha = \bigcup_n [f_\alpha^{2n}(0), f_\alpha^{2n+1}(0))$ and define $\mathcal{F} = \langle X_\alpha : \alpha < \mathfrak{b} = \mathfrak{d} \rangle$. We must show that $\mathcal{F}$ is a non-meager P^+ -filter.

Lemma 3.7 $(\forall A \in [\omega]^\omega)(\exists \alpha)(\forall^\infty n) A \cap [f_\alpha^n(0), f_\alpha^{n+1}(0)) \neq \emptyset$.

Proof: Given A , define $f \in {}^\omega\omega$ such that $A \cap [n, f(n)) \neq \emptyset$ for each n . Now there must be an α such that $f \leq^* f_\alpha$ and therefore

$$(\forall^\infty n) A \cap [n, f_\alpha(n)) \neq \emptyset.$$

Thus $(\forall^\infty n) A \cap [f_\alpha(n), f_\alpha(f_\alpha(n))) \neq \emptyset$, and finally

$$(\forall^\infty n) A \cap [f_\alpha^n(0), f_\alpha^{n+1}(0)) \neq \emptyset$$

as desired. $\square$

We are ready to show that $\mathcal{F}$ is a P^+ -filter. Fix a descending sequence $Y_0 \supseteq Y_1 \supseteq \dots$ from $\mathcal{F}^+$. By induction, define a sequence of ordinals $\alpha_n < \mathfrak{b} = \mathfrak{d}$ by $\alpha_0 = 0$ and given α_n , let $\mathcal{F}_n$ be the filter generated by

$$\{X_\gamma : \gamma < \alpha_n\} \cup \{Y_k : k \in \omega\}.$$

Now choose $\alpha_{n+1} > \alpha_n$ so that

$$(\forall Y \in \mathcal{F}_n)(\forall^\infty k) Y \cap [f_{\alpha_{n+1}}^k(0), f_{\alpha_{n+1}}^{k+1}(0)) \neq \emptyset.$$

This is possible using the lemma and the fact that $\mathcal{F}_n$ is generated by less than $\mathfrak{b} = \mathfrak{d}$ sets. Finally let $\alpha = \sup\{\alpha_n : n \in \omega\}$. We thus get

$$(\forall Y \in \mathcal{F}_\alpha := \langle X_\gamma : \gamma < \alpha \rangle \cup \langle Y_n : n \in \omega \rangle)(\forall^\infty k) Y \cap [f_\alpha^k(0), f_\alpha^{k+1}(0)) \neq \emptyset.$$

Therefore it suffices to find $Z \subseteq^* Y_n$ such that

1. $Z \in \mathcal{F}_\alpha^+$
2. $(\forall^\infty k) Z \cap [f_\alpha^k(0), f_\alpha^{k+1}(0)) \neq \emptyset$.

But again, $\mathcal{F}_\alpha$ is generated by less than $\mathfrak{b}$ sets, say by $\langle X_\beta : \beta < \lambda \rangle$ for some $\lambda < \mathfrak{b}$, and we may assume that this collection is also closed under finite intersections. Now for each such β , choose $h_\beta \in {}^\omega\omega$ such that $Z_\beta = \bigcup_n Y_n \cap h_\beta(n)$ satisfies:

1. $(\forall^\infty k) Z_\beta \cap [f_\beta^k(0), f_\beta^{k+1}(0)) \neq \emptyset$.
2. $Z_\beta \cap X_\beta$ is infinite.

Now choose $h^* > h_\beta$ for $\beta < \lambda$. Then $Z = \bigcup_n Y_n \cap h^*(n)$ is as desired.

Example 3.8 (CH) A strong P^+ -filter not included in any F_σ filter.

Enumerate all closed sets which generate a proper filter as $\langle \mathcal{C}_\alpha : \alpha < \omega_1 \rangle$ and all sequences of potential candidates for members of $\mathcal{F}^+$ as $\langle X_n^\alpha : n \in \omega \rangle$ for $\alpha < \omega_1$. We build a filter $\mathcal{F}$ in ω_1 stages. At stage α , assume that we have built for $\beta < \alpha$:

1. $\langle a_n^\beta : n \in \omega \rangle$ a sequence of finite sets.
2. $X_\beta \in [\omega]^\omega$ such that $\lim_n |X_\beta \cap a_n^\gamma| = +\infty$ for all $\gamma < \alpha$.
3. $\beta < \gamma < \alpha \rightarrow X_\gamma \subseteq^* X_\beta$.

Let $\mathcal{F}_\alpha$ be the filter generated by $\{X^c : (\forall n)|X \cap a_n^\beta| \leq m\}$ for $m \in \omega$ and by $\{X_\beta : \beta < \alpha\}$. We are given a closed set $\mathcal{C} = \mathcal{C}_\alpha$ and we want to make sure that $\mathcal{F}_\alpha \not\subseteq \langle \mathcal{C} \rangle$. So we suppose that in fact that $\mathcal{F}_\alpha \subseteq \langle \mathcal{C} \rangle$ and we may as well assume that $\alpha = \omega$.

Choose a sequence $n_0 < n_1 < \dots$ such that

$$(\forall Y_1, \dots, Y_k \in \mathcal{C}) Y_1 \cap \dots \cap Y_k \cap [n_k, n_{k+1}) \neq \emptyset.$$

Claim: $(\forall \beta < \alpha)(\forall m)(\forall Z)$, if $|Z \cap a_k^\beta| \leq m$ for each k , then

$$(\forall Y \in \mathcal{C})(\forall^\infty \ell) (Y \cap [n_\ell, n_{\ell+1})) \setminus Z \neq \emptyset.$$

Proof: Indeed we have made sure to put Z^c in $\mathcal{F}_\alpha$ for all such Z 's, and $\mathcal{F}_\alpha \subseteq \langle \mathcal{C} \rangle$ by assumptions. $\square$

The construction of X_α can therefore be obtained by a dovetailed construction. Now to handle the sequence $\langle X_n^\alpha : n \in \omega \rangle$, given that each $X_n^\alpha \in \langle \mathcal{F}_\alpha, X_\alpha \rangle^+$, choose $f \in {}^\omega \omega$ such that

$$(\forall n) |X_\alpha \cap X_n^\alpha \cap [n, f(n))| \geq n,$$

and define $a_n^\alpha = [n, f(n)) \cap X_n^\alpha$.

This completes the construction. $\mathcal{F} = \bigcup_\alpha \mathcal{F}_\alpha$ is the desired filter.

Example 3.9 (CH) A mad family $\mathcal{A}$ such that $\mathcal{F}(\mathcal{A})$ is a strong P^+ -filter.

In the following, $\mathcal{I}(\mathcal{A})$ denotes the ideal generated by members of $\mathcal{A}$. We start with a lemma.

Lemma 3.10 (CH) There is a mad family $\mathcal{A}$ such that

$$(\forall X \notin \mathcal{F}(\mathcal{A}) \cup \mathcal{I}(\mathcal{A}))(\exists \alpha)(\forall \beta > \alpha) A_\beta \setminus X \text{ and } A_\beta \cap X \text{ are infinite}.$$

Proof: Start with a partition of ω into infinite sets $\langle A_n : n \in \omega \rangle$. Now enumerate $[\omega]^\omega$ as $\langle X_\alpha : \alpha < \omega_1 \rangle$. Suppose that at stage $\alpha < \omega_1$, we have $\mathcal{A}_\alpha = \langle A_\beta : \beta < \alpha \rangle$ and $\mathcal{X}_\alpha \subseteq [\omega]^\omega \setminus [\mathcal{F}(\mathcal{A}_\alpha) \cup \mathcal{I}(\mathcal{A}_\alpha)]$. Relist $\mathcal{A}_\alpha$ as $\langle B_n : n \in \omega \rangle$.

Case 1: $X_\alpha \in \mathcal{F}(\mathcal{A}_\alpha) \cup \mathcal{I}(\mathcal{A}_\alpha)$. There is nothing to do.

Case 2: $X_\alpha \notin \mathcal{F}(\mathcal{A}_\alpha) \cup \mathcal{I}(\mathcal{A}_\alpha)$. Thus in particular both $X_\alpha \not\subseteq^* B_0 \cup \dots \cup B_n$ and $X_\alpha^c \not\subseteq^* B_0 \cup \dots \cup B_n$ for each n .

List $\mathcal{X}_\alpha \cup \{X_\alpha\}$ as $\langle X_n : n \in \omega \rangle$ and we build A_α such that:

1. $(\forall n) A_\alpha \cap B_n =^* \emptyset$.

2. $(\forall n) A_\alpha \setminus X_n$ and $A_\alpha \cap X_n$ are infinite .

At stage n , assume that we have $a_n \in [\omega]^{<\omega}$, then choose

$$k_i \in X_i \setminus (B_0 \cup \dots \cup B_n \cup n) \text{ for } i < n,$$

$$\ell_i \in X_i^c \setminus (B_0 \cup \dots \cup B_n \cup n) \text{ for } i < n,$$

and let $a_{n+1} = a_n \cup \{k_i, \ell_i\}_{i < n}$. Hence $A_\alpha = \bigcup_n a_n$ satisfies 1-2 and put $\mathcal{A}_{\alpha+1} = \mathcal{A}_\alpha \cup \{A_\alpha\}$, $\mathcal{X}_{\alpha+1} = \mathcal{X}_\alpha \cup \{X_\alpha\} \subseteq [\omega]^\omega \setminus [\mathcal{F}(\mathcal{A}_{\alpha+1}) \cup \mathcal{I}(\mathcal{A}_{\alpha+1})]$. This completes the construction. The verification that $\mathcal{A} = \{A_\alpha : \alpha < \omega_1\}$ has the desired properties is straightforward. $\square$

Now we show that such a mad family yields a strong P^+ -filter $\mathcal{F}(\mathcal{A})$. But given a sequence $\langle X_n : n \in \omega \rangle$ from $\mathcal{F}(\mathcal{A})^+$, we can choose by the assumptions on $\mathcal{A}$ countably members $\langle A_n : n \in \omega \rangle$ from $\mathcal{A}$ such that $A_n \cap X_m$ is infinite for each n and m . Now define $f \in {}^\omega \omega$ such that

$$(\forall n)(\forall i < n) A_i \cap X_n \cap [n, f(n)) \neq \emptyset.$$

If $Y \in \mathcal{F}(\mathcal{A})$, there must be an A_n such that $A_n \subseteq^* Y$ and therefore $Y \cap X_n \cap [n, f(n)) \neq \emptyset$ for all but finitely many n .

4 Conclusion

The two implications that remain to be solved are the following.

Question 1: Does a completely meager P^+ -filter, or actually even a hereditarily meager filter necessarily have the strong Baire property?

One remains in ZFC.

Question 2: Is there a strong P^+ -filter not included in any F_σ -filter?

We still have very little understanding of the structure of filters of the form $\mathcal{F}(\mathcal{A})$ for a mad family $\mathcal{A}$. In particular, the following remains unsolved.

Question 3: Do all filters of the form $\mathcal{F}(\mathcal{A})$ for a mad family $\mathcal{A}$ have the strong Baire property?

We however have a meagerness property that separates filters of the form $\mathcal{F}(\mathcal{A})$ and F_σ -filters, namely the strong P^+ property. In view of the results of [3] that F_σ -filters can be diagonalized in a forcing extension without adding an unbounded real, we may ask the following.

Question 4: Can all strong P^+ -filters be diagonalized in a forcing extension without adding unbounded reals?

Of course the fundamental motivation for these questions is:

Question 5: Can filters of the form $\mathcal{F}(\mathcal{A})$ for a mad family $\mathcal{A}$ be diagonalized in a forcing extension without adding an unbounded real?

References

- [1] T. Bartoszynski and H. Judah, *Measure and Category - Filters on ω* , Set Theory and the Continuum, H. Judah, W. Just and H. Woodin eds., (1992), Springer-Verlag.
- [2] W. Just, A.R.D. Mathias, K. Prikry and P. Simon, *On the existence of large P -ideals*, to appear.
- [3] C. Laflamme, *Zapping small filters*, Proc. Amer. Math. Soc. 114 (1992), 535-544.
- [4] S. Shelah, *Cardinal invariants of the continuum*, Axiomatic Set Theory, J. Baumgartner, D. Martin and S. Shelah eds., Contemporary Mathematicsvol. 31, AMS 1984.
- [5] P. Simon, private communication, August 1987.
- [6] M. Talagrand, *Compacts de fonctions mesurables et filtres non mesurables*, Studia Math. 67 (1980), 13-43.
- [7] M. Talagrand, *Filtres: mesurabilité, rapidité, propriété de Baire forte*, Studia Math. 74 (1982), 283-291.